

Subcutaneous Injections of Nanofat Adipose-derived Stem Cell Grafting in Facial Rejuvenation

Sophie Menkes, MD*
 Mariotta Luca, PhD†
 Gianni Soldati, PhD†
 Luigi Polla, PhD*

Background: We aimed to assess whether our novel Nanofat grafting procedure improves skin quality while yielding a regenerative effect and whether this novel technique can also achieve a lifting effect.

Methods: Patients who requested nonsurgical facial rejuvenation were enrolled between June 2018 and December 2018. Fat was aspirated from the medial thigh, inner part of the knee, or lower abdomen regions. Following aspiration and flushing, microfat was obtained after washing with saline. This microfat was emulsified to obtain a Nanofat suspension, which was injected using a 25-G cannula into the subcutaneous layer at different facial sites. Images were obtained before and at 1, 3, and 6 months after facial rejuvenation. Patients were also administered a survey. Characterization of the isolated stromal vascular fraction (3 patients), and before/after biopsies were performed.

Results: Fifty patients were included (2 men and 48 women; mean age, 35–65 years; mean follow-up, 9 months). The clinical results were apparent between 2 and 4 weeks after injection, and improvements were continuously observed until 6 months postoperatively. All patients confirmed an improvement in skin quality. A lifting effect was also observed. The data confirm that the Nanofat procedure does not damage cells, maintaining cell viability, and number of adipose-derived stem cells. Biopsies showed an increased dermal cellularity, vascular density, and elastic and collagen fiber density.

Conclusion: Facial rejuvenation with subcutaneous Nanofat injections appears to be an effective method, representing a skin rejuvenation effect by modifying the pattern of the dermis, although additional studies are necessary. (*Plast Reconstr Surg Glob Open* 2020;8:e2550; doi: [10.1097/GOX.0000000000002550](https://doi.org/10.1097/GOX.0000000000002550); Published online 21 January 2020.)

INTRODUCTION

The demand for nonsurgical facial restoration has been constantly increasing in the past few decades. The Nanofat grafting procedure is a promising technique that could meet this demand and yields a marked improvement in skin quality. (See **Video [online]**, which displays facelift by Nanofat grafting.)

In 2013, Tonnard et al introduced a Nanofat grafting technique, wherein mechanical emulsification was used to mechanically dissociate adipose tissue while retaining the extracellular matrix with all nucleated cells including the already present stem cells. This sample was then injected

using a 30-G needle into the fine lines of the face.¹ This method took advantage of the release of growth factors from cells present in the tissue and the regenerative capacities of adipose-derived stem cells (ASCs).^{2,3} On exposure to tissue-specific biochemical signals, ASCs could differentiate into different tissues. This grafting procedure frees the cells present in the adipose tissue to have a biologic effect in the process of reshaping the face without a change in volume.

In the present study, we aimed to assess whether the Nanofat grafting procedure improves skin quality and whether the product issued from this process of mechanical dissociation maintains the cells present in the tissue alive and in a reasonable amount. Finally, we also sought to determine whether this novel technique can also achieve a lifting effect.

MATERIALS AND METHODS

Fat Collection

We enrolled patients who requested nonsurgical facial rejuvenation between June 2018 and December 2018. The

From the *Forever Institut, Geneva, Switzerland; and †Swiss Stem Cell Foundation in Pasquée, Gentilino, Switzerland.

Received for publication July 24, 2019; accepted September 4, 2019.

Copyright © 2020 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: [10.1097/GOX.0000000000002550](https://doi.org/10.1097/GOX.0000000000002550)

Disclosure: The authors have no financial interest to declare in relation to the content of this article.

Nanofat grafting procedure was performed under local anesthesia. In brief, following disinfection with povidone iodine, fat was aspirated at the most superficial level possible to recover the maximum number of stem cells from the medial thigh, inner part of the knee, or lower abdomen. Anesthesia was induced with infiltration with Klein solution (500-mL normal saline, 50-mL 2% lidocaine, 1-mL 1:1,000 adrenaline, and 15-mL 8.4% sodium bicarbonate) between 1 and 4 areas are injected with 20–40 mL. For fat collection, we used the Hapi Fat kit (Benewmedical, Geneva, Switzerland); a 14-G 130-mm cannula was connected to a 10-mL Luer Lock syringe, and small adipocyte lobules were harvested under low negative pressure via a fatlock connector valve system and a tube attached to a PureGraft 50 pocket.

Pure Graft Procedure

The procedure was conducted in a closed system, and the quantity of fat harvested ranged from 30 to 50 mL. Each 10-mL sample of fat aspirated and flushed into the Pure Graft pocket was washed with 10 mL of sterile saline. At the end of the procedure, 40 mL of fat was collected to obtain approximately 20 mL of microfat tissue.

Emulsification of the microfat to obtain a Nanofat suspension was achieved with the Tulip kit (Tulip Medical products, San Diego, CA); this included 3 filters (2.4, 1.4, and 1.2 mm) and involved shifting between two 20-mL syringes connected to each other by a female-to-female Luer-lock connector for >30 passes for each filter. The final filter included a Tulip nanotransfer 600/400 μm device, and only 1 pass was used to obtain a pure product with the least number of fibers possible; the product could easily be injected through a 25-G cannula or a 27-G needle.

A total of 15 mL of Nanofat suspension was expected to be recovered from 20 mL of microfat. To this suspension, we added 20% platelet-rich plasma (PRP), which was prepared from the native blood of the patient with a 2-step centrifugation Duografter II (Proteal, Madrid, Spain), after the platelets were counted with a Horiba ABS Micro device (Axonlab, Baden, Switzerland) to ensure appropriate quality for Nanofat grafting.

The Nanofat-PRP suspension was finally transferred into individual 1-mL syringes and could be injected into the recipient area. The emulsion was injected with a fanning technique, using a 25-G cannula into the subcutaneous layer at various sites on the face via 4 entry points: orbital area, cheek, jaw line, and temporal area. Nanofat grafting was first performed at the lower lid and orbital area 1 mL, followed by the upper lip 1 mL, lower lip 1 mL, chin 1 mL, cheek 1 mL, nasolabial fold 1 mL, jaw line 1 mL, parotid area 1 mL, and temporal area 1 mL. Nanofat was applied from a single syringe in each area (Fig. 1).

Standardized photographs were obtained for all patients before the procedure, and at 1, 3, and 6 months after facial rejuvenation. All the patients were self-evaluated using a visual scale, graded from 0 to 10. The survey concerned the following clinical aspects of the face: luminosity, firmness, superficial and deep wrinkles, folds, volume, and overall rejuvenation. The subjects were also

asked to grade the overall appreciation by their social circle (Fig. 2).

Characterization of the Isolated Stromal Vascular Fraction

The fraction produced by the Nanofat procedure was characterized for cell counting, viability, and ASC presence by enzymatically extracting the Nanofat product. The sample was washed with 40-mL Dulbecco's Deprievied Phosphate Buffered Saline (DPBS; with Ca^{2+} and Mg^{2+} ; PAA Laboratories, Pasching, Austria) by gentle agitation. The syringe was held vertically in the support stand for a few minutes to allow for the separation of the phases, then the lower aqueous phase was discarded by pushing the piston. The sample was washed twice. To free the cells in the aqueous phase, the washed adipose tissue must be digested with the appropriate amount of Liberase MTF-S (Roche Applied Science, Basel, Switzerland) at a final concentration of 0.28 Wünsch U/mL diluted in 10-mL DPBS (with Ca^{2+} and Mg^{2+}). The sample was incubated for 45 minutes at 37°C under constant but gentle agitation. Enzymatic reaction was stopped by aspiration of 30 mL of injectable 5% human albumin solution (CSL Behring AG, Bern, Switzerland) in the syringe. The syringe was then put back in vertical position to allow the separation of the phases. The lower layer, which now contained the stromal vascular fraction (SVF) cells, was carefully poured out into a conical 50-mL centrifuge tube (TPP, Trasadingen, Switzerland). The extracted cells were washed again with 40-mL 5% human albumin solution to increase cell yield. Finally, after filtration through a 100- and 40- μm sieve (Cell Strainer, BD Falcon, Basel, Switzerland), SVF was centrifuged 400g, 5-minute room temperature (RT) and the pellet suspended another time in DPBS (without Ca^{2+} and Mg^{2+} ; PAA Laboratories) or in tissue culture medium. The SVF was then analyzed for cell count and viability using an automated propidium iodide-based cell counting device (Nucleocounter NC-100, Chemometec A/S, Denmark). The cells of the SVF were also characterized by cytofluorimetric analysis using a 10-channel Navios cytometer [Beckman Coulter (BC), Nyon, Switzerland]. Briefly, roughly 500,000 cells from fresh SVF preparation were taken and centrifuged 5 minutes at 400g. The pellet was resuspended in 220 μL of PBS without Ca^{2+} / Mg^{2+} (Eurobio, CS1PBS01) with 1% human converted AB serum (PAA, C11-021), and the cells were stained for

Nanofat grafting in subcutaneous layer: an innovative technique




Fig. 1. Graph of injection sides.



Fig. 3. Photographs of the face before (A) and 6 months after (B) 18-mL Nanofat grafting of a 42-year-old patient.



Fig. 4. Photographs of the face before (A) and 6 months after (B) 18-mL Nanofat grafting of a 48-year-old patient.

achieved with the procedure. In the survey (Fig. 7), most patients answered that they now did not need to apply makeup or skin care daily. All patients confirmed an improvement in texture, elasticity, glow, firmness, fine wrinkles, and skin hydration, along with a regenerative effect. Moreover, all patients exhibited a considerable improvement with regard to skin glow and regularity, and more than 80% of patients exhibited a large improvement in facial shape and opinions of relatives.

No major side effects were observed in the present study. Minor complications, including redness and edema between 2 and 4 days, some bruises, and pain at the donor sites, were noted in all the patients. Only 1 patient developed nodules at lower lid, although this condition healed spontaneously after 1 month. No infections, fat cysts, or granulomas were observed after treatment and during the follow-up period.

To assess the quality of Nanofat product, we extracted the SVF of 3 samples after Nanofat processing. Data are




Fig. 5. Photographs of the face before (A) and 6 months after (B) 18-mL Nanofat grafting of a 52-year-old patient.




Fig. 6. Photographs of the face before (A) and after (B) 6 months 18-mL Nanofat grafting of a 35-year-old patient.

shown in Fig. 8. As can be observed, the viability of the cells is maintained after this procedure of mechanical disruption with more than 90% of cell viable. Total nucleated cells counts are also in the range of what is obtained by regular SVF extraction without Nanofat procedure, where, on more than 300 samples processed under GMP guidelines, we found a comparable mean value. Finally, the number of ASCs is also similar to that obtained without the Nanofat procedure.

Overall, the data confirm that the Nanofat procedure does not damage cells contained in the adipose tissue, maintaining cell viability, and number of ASCs.

Histologic analyses were performed to assess the histologic changes of the skin after injection of Nanofat grafting.

Skin biopsy performed on the posttreatment zone showed an increase in collagen (denser and broader collagen fibers) and elastic fibers (control: 171 ± 9.5 ; Nanofat: 266 ± 17.3 ; $P < 0.005$) in the superficial dermis (Fig. 9). Treated zones also displayed an increased dermal cellularity (mainly fibroblasts) (control: 59.6 ± 6.8 ; Nanofat: 90.6 ± 3.05 ; $P < 0.03$) and vascular density (control: 15.6 ± 0.5 ; Nanofat: 26.6 ± 3.05 ; $P < 0.01$) in the superficial dermis (Fig. 10). No modification was observed in epidermal thickness or in mucin content of the dermis. There was no significant inflammation in the treated zones.

DISCUSSION

The quality of an individual's skin communicates several details about their habits. There are 2 major factors that form the foundation of patients' attitudes toward skin quality: health (radiant, glow, natural, nourished) and youth (no wrinkles, smooth, firm, plump, elastic). Patients believe that skin beauty plays a role in self-confidence and success.

The Nanofat grafting technique seeks to increase vascularization, neocollagenesis, and tissue regeneration and also to improve skin physiology. In particular, ASCs play a vital role in skin rejuvenation,² although their multipotent nature may not be the only advantage. ASCs also secrete stimulatory factors, such as cytokines, chemotactic factors, extracellular remodelers, growth factors, and their inhibitors. These factors are capable of very




Fig. 7. Patients' answers to questions about skin improvement.




Fig. 8. Comparison of N = 3 patients, where adipose samples have been extracted before and after Nanofat processing. In both cases, cells were extracted by manual processing following enzymatic digestion and then counted for total nucleated cells and ASCs.




Fig. 9. Hematoxylin–eosin staining (A, B) and CD31 immunohistochemistry (C, D) of untreated (A, C) and Nanofat-treated (B, D) skin (original magnification, 20×). Note the increased density of dermal cellularity (mainly fibroblasts) (B) and dermal vessels (D) in the treated skin.

complex functions involving several bioactive factors and reactivities.³ ASCs not only help in skin regeneration but also in the repair of various functions such as melanocyte action. ASCs inhibit melanocyte proliferation and melanin synthesis by downregulating melanogenic enzymes through an interleukin-6-mediated mechanism.^{4,5} This

leads to good results for the treatment of dark coloration of lower eyelid skin.⁶

In a previous study, Tonnard et al quantified the cells from the SVF and its CD34+ subfraction; the researchers assessed stem cell quality by culturing the cells in standard and adipogenic media.¹ In the emulsion, the normal fat




Fig. 10. Trichrome-Masson (A, B) and Miller (C, D) stainings of untreated (A, C) and Nanofat-treated (B, D) skin (original magnification, 20 \times). Note the increased density of collagen (B) and elastic fibers (D) in the treated skin.

tissue structure is lost, but mesenchymal stem cells remain intact as they were in the undissociated adipose tissue.⁷ In the 3 samples analyzed, we found higher number of ASCs than microfat samples (Fig. 7). Moreover, these cells are associated with increased regenerative capacity, which may have contributed to the observed lifting effect.⁸ Nanofat processing is more than simply breaking down adipocytes; in fact, mechanical emulsification leads to upregulation of multipotent and pluripotent markers, which induce the regenerative capacity.¹ In addition to skin regeneration, the product also created a lifting effect through the injection of the product subdermally in many areas of the face. This technique led to improvements in facial appearance without a change of volume.¹

Hyaluronic acid has been used as an appropriate treatment for improving skin quality. In addition, lipofilling, PRP, mesotherapy with peptides and vitamins, lasers, and laser-based devices such as ultrasound and radiofrequency are also considered appropriate options. However, these modalities only have a moderate impact on skin quality, and fillers have also been reported to cause complications, lumps, bruising, and irregularities.⁹

Additional studies should be conducted to test a control group with hyaluronic acid or PRP alone. Liang et al showed that both Nanofat-plasma rich fibrin (PRF) and hyaluronic acid injection improve facial skin status without serious complications with greater patient satisfaction, implying that Nanofat-PRF injection is a safe, highly effective, and long-lasting method for skin rejuvenation.¹⁰

We believe that the lower third of the aging face (lips and immediate perioral area) deserves as much attention as the midface and that the temporal area should be covered with the injections. The injection of Nanofat in the subcutaneous layer could enable expansion of this intervention to sensitive areas, including skin rejuvenation

from the zygomatic arch to the temporal area, without side effects.

In the Nanofat procedure, the time required for infiltration, harvesting, and emulsification ranged from 30 to 45 minutes; to reduce process time, PRP was centrifuged before fat harvesting. The injection time with the cannula remained the same as for treatment with fillers. The Nanofat grafting procedure may cause swelling for 2–4 days, as well as bruising and pain at the donor site for 1 week. However, no delayed reactions have been noted with this procedure over 5 years, as noted through studies in the literature. To avoid vascular complications, the product was injected with a cannula; if a needle was used, small aliquots would need to be injected under low pressure.

The disadvantages of the Nanofat technique include the necessity of a surgical procedure, as some discomfort is involved in extracting fat from the donor site. Moreover, the site and manner in which the product is injected are important factors for obtaining a good result, but there are other factors involved as well. Each step in the process should be carefully monitored, beginning with the choice of donor site, followed by harvesting, processing, and injection techniques.^{11–14} Disinfection is very important, although Pallua et al showed that local antiseptic treatment can affect ASC viability. In particular, octenidine has been found to decrease ASC viability, even at the lowest concentrations.¹⁵ Povidone iodine and polyhexanide also decrease ASC viability and significantly reduce the expression of stem cell markers (CD29, CD34, CD73, CD90, CD105). Polyhexanide reduces the levels of CD34 alone, but mafenide does not alter ASC viability, even at high concentrations, and does not affect stem cell marker levels. Thus, mafenide is a good antiseptic for such purposes. However, in the present study, we used povidone iodine as it was easy to procure and then

washed the area carefully with saline solution, including the entry points, to avoid any introduction of antiseptic inside the tissues.¹⁵

Furthermore, Pallua et al indicated that the abdominal wall is the optimal site for harvesting due to the high number of SVF cells, as well as high levels of MMP-9, IGF-1bFGF, and PDGF. However, there was no significant difference in the ASC count (CD90+, CD105+, CD31-) among the harvest sites, and the proliferation, differentiation, and SVF cell/ASC counts did not significantly differ among the harvest sites.¹⁶ In addition, the amount of anesthesia administered should be limited to protect the stem cells, as lidocaine has been found to be toxic.¹⁷

Tonnard et al indicated that microfat should be harvested with barbed (rasping) cannulas, as they contain a greater number of ASCs as compared to fat harvested with smooth multiholed cannulas.¹² In the present study, we used a microcannula as it represented a minimally invasive method to obtain clinically relevant numbers of stromal and vascular cells for tissue regeneration and neovascularization.¹⁸ Furthermore, the collected material had to be clear and washed carefully. Condé-Green et al observed that washed fat grafts contained higher amounts of ASCs, as compared to decanted and centrifuged fat grafts.¹⁹ By adding PRP, we enhance cell graft survival and maintenance.^{20,21}

PRP alone is effective on skin rejuvenation, but a minimum of 3 sessions are still necessary to obtain results.²² In a thesis work on PRP as potentialiser of cell therapy in the field of plastic surgery, 20% is the number allowing the best efficiency.²³

Some data suggest that lesions could be treated by injection of ASCs. They are capable to differentiate in other cells giving them the first role in regenerative medicine. They also have an important paracrine activity and modulate immune tolerance and inflammation.²⁴ The effects of Nanofat are visible in the epidermis also if the injection is in the subcutaneous layer. We need however more studies to understand all sequences of events. It is interesting to note that treatment with Nanofat induces histologic modification quite similar to those induced by SVF. It modifies the pattern of collagen fibers and elastic fibers in the dermis.

Biopsy was performed in the inner part of the forearm because biopsy on the face was not acceptable for patients. The skin is not similar in all parts of the body, even in the face. Distribution of skin thickness (diameter of the epidermis and dermis) differs in the face from 2.5 mm in the chin to 0.5 mm in the eyelids. The number of hair follicles per square centimeter differs too, from 700 on the chin to 1,200 on the nose. But the layers, the structure, function, and number of fibroblasts are the same in all the skin types.²⁵

Fat grafting techniques are generally based on the volume and goal needed, including the Coleman technique [small volume fat grafting (<150 mL) for facial rejuvenation and regeneration], Khouri technique [large-volume fat grafting (>150 mL) for breast contouring and buttock augmentation], and Nanofat technique by Tonnard [very small volume (<15 mL) for facial skin rejuvenation with a small and sharp needle for intradermal injections).

At the Forever Institut (Geneva, Switzerland), we developed a novel technique wherein Nanofat (18 mL) was subcutaneously injected at many facial sites using a 25-G cannula, and good results were noted in terms of skin quality and a lifting effect. The duration for which the effects of Nanofat grafting persist is unclear, as this procedure needs to be assessed further. Hence, additional studies should be conducted to test the significant effect of Nanofat grafting¹ and should include objective skin quality tests with a Cutometer, to measure improvement, elasticity, and firmness.

CONCLUSIONS

Although the current modalities are all suitable for improving skin quality, facial rejuvenation with subcutaneous Nanofat injections appears to be a very effective method, by modifying the pattern of dermis. Nanofat grafting is a regenerative technique, involves an autologous procedure, enables easy collection, and involves single-day ambulatory treatment. In addition, the results of clinical trials indicated the safety of the method. Moreover, Nanofat naturally integrates into host tissues without producing any major side effects. The long-term effects need to be further evaluated, but Nanofat technology can likely be expanded to cover indications for hair or genital rejuvenation; the initial results for those indications appear very promising.

Sophie Menkes, MD

Forever Institut

56 rue du Rhône

1204 Geneva, Switzerland

Email: smenkes@forever-beauty.com

REFERENCES

1. Tonnard P, Verpaele A, Peeters G, et al. Nanofat grafting: basic research and clinical applications. *Plast Reconstr Surg*. 2013;132:1017–1026.
2. Charles-de-Sa L, Gontijo-de-Amorim NF, Maeda Takiya C, et al. Antiaging treatment of the facial skin by fat graft and adipose-derived stem cells. *Plast Reconstr Surg*. 2015;135:999–1009.
3. Caplan A, Yap S, Casteilla L, et al. Perivascular origin for mesenchymal stem cells in multiple human organs. *Cell Stem Cell*. 2008;3:301–313.
4. Lin JY, Fisher DE. Melanocyte biology and skin pigmentation. *Nature*. 2007;445:843–850.
5. Deok-Woo K, Byung Joon J, Na Hyun H, et al. Adipose derived stem cells inhibit epidermal melanocytes through an interleukin-6-mediated mechanism. *Plast Reconstr Surg*. 2014;134:470–480.
6. Dong Seok O, Dae Hwa K, Tai Suk R, et al. Correction of dark coloration of the lower eyelid skin with Nanofat grafting. *Arch Aesthetic Plast Surg*. 2014;20:92–96.
7. Memar O, Nezamabadi A, Milani BY, et al. Nanofat grafting: basic research and clinical application. *Plast Reconstr Surg*. 2014;133:728e.
8. Banyard DA, Sarantopoulos CN, Borovikova AA, et al. Phenotypic analysis of stromal vascular fraction after mechanical shear reveals stress-induced progenitor populations. *Plast Reconstr Surg*. 2016;138:237e–247e.
9. Goisis M, Stella E, Di Petrillo A, et al. Nanofat grafting compared to hyaluronic acid and PRP treatment of the lower lid and tear trough: clinical outcome. *JSM Ophthalmol*. 2015;3:1057.
10. Liang ZJ, Lu X, Li DQ, et al. Precise intradermal injection of nanofat-derived stromal cells combined with platelet-rich fibrin

- improves the efficacy of facial skin rejuvenation. *Cell Physiol Biochem*. 2018;47:316–329.
11. Alharbi Z, Opländer C, Almakadi S, et al. Conventional vs. micro-fat harvesting: how fat harvesting technique affects tissue-engineering approaches using adipose tissue-derived stem/stromal cells. *J Plast Reconstr Aesthet Surg*. 2013;66:1271–1278.
 12. Caggiati A, Germani A, Di Carlo A, et al. Naturally adipose stromal cell-enriched fat graft: comparative polychromatic flow cytometry study of fat harvested by barbed or blunt multihole cannula. *Aesthet Surg J*. 2017;37:591–602.
 13. Charles-de-Sá L, Gontijo-de-Amorim NF, Maeda Takiya C, et al. Antiaging treatment of the facial skin by fat graft and adipose derived stem cells. *Plast Reconstr Surg*. 2015;135:999–1009.
 14. Rodriguez RL, Condé-Green A. Reply: adipocyte damage in relation to different pressures generated during manual lipoaspiration with a syringe. *Plast Reconstr Surg*. 2013;131:646e–647e.
 15. Kim BS, Ott V, Boecker AH, et al. The effect of antiseptics on adipose-derived stem cells. *Plast Reconstr Surg*. 2017;139:625–637.
 16. Kakagia D, Pallua N. Autologous fat grafting: in search of the optimal technique. *Surg Innov*. 2014;21:327–336.
 17. Wu T, Smith J, Nie H, et al. Cytotoxicity of local anesthetics in mesenchymal stem cells. *Am J Phys Med Rehabil*. 2018;97:50–55.
 18. Trivisonno A, Di Rocco G, Cannistra C, et al. Harvest of superficial layers of fat with a microcannula and isolation of adipose tissue-derived stromal and vascular cells. *Aesthet Surg J*. 2014;34:601–613.
 19. Condé-Green A, de Amorim NF, Pitanguy I. Influence of decantation, washing and centrifugation on adipocyte and mesenchymal stem cell content of aspirated adipose tissue: a comparative study. *J Plast Reconstr Aesthet Surg*. 2010;63:1375–1381.
 20. Liao HT, James IB, Marra KG, et al. The effects of platelet-rich plasma on cell proliferation and adipogenic potential of adipose-derived stem cells. *Tissue Eng Part A*. 2015;21:2714–2722.
 21. Jin R, Zhang L, Zhang YG. Does platelet-rich plasma enhance the survival of grafted fat? An update review. *Int J Clin Exp Med*. 2013;6:252–258.
 22. Fisher GJ, Wang ZQ, Datta SC, et al. Pathophysiology of premature skin aging induced by ultraviolet light. *N Engl J Med*. 1997;337:1419–1428.
 23. Hersant B. *Role of PRP as potentiator of cell therapy in the field of plastic surgery [unpublished thesis]*.
 24. Lee SH, Jin SY, Song JS, et al. Paracrine effects of adipose-derived stem cells on keratinocytes and dermal fibroblasts. *Ann Dermatol*. 2012;24:136–143.
 25. Rubin MG, Schurer NY, Wiest LG, et al. *Illustrated guide to chemical peels: basics indications, uses*. Batavia, IL: Quintessence Publishing Group.